arXiv:1208.2961v1 [physics.gen-ph] 15 Aug 2012

Why the concept of frozen-in field lines is often misleading

Zhu-Xing Liang[1, a)] and Yi Liang[1]
18-4-102 Shuixiehuadu, Zhufengdajie, Shijiazhuang, Hebei 050035, China

(Dated: 22 October 2012)

According to the concept of frozen-in field lines, a highly conducting fluid can move freely along magnetic field lines but cannot freely traverse them. Here we show analytically that under some special conditions, the frozen-in field concept may not hold. Under some conditions, whether a conducting fluid can freely cross magnetic field lines or not appears to depend solely on the gradient of the magnetic field in the direction of fluid movement and is independent of the magnetic field strength. We especially emphasize the significance of isomagnetic surfaces and polarization charges. We also present experimental results to support our analysis.

PACS numbers: 52.30.Cv, 95.30.Qd, 94.05.Rx, 94.30.cs
Keywords: magnetohydrodynamics, plasma, experiment

I. INTRODUCTION

The physical properties of magnetohydrodynamics (MHD) – the frozen-in property of field lines and the conservation of magnetic flux[2] – are often referred to collectively as Alfvén's Theorem. The concept of frozen-in field lines has been presented in various forms. A common interpretation is that a highly conducting fluid can move freely along, but cannot freely traverse, magnetic field lines. Because the frozen-in concept is of prime importance to plasma physics, it is widely mentioned in the literature,[12–16] and is presented in almost all textbooks of plasma physics.[7,8] Although the applicability of this concept has long been contested (indeed,[1,13] and Alfvén, the concept's founder, criticized its use and stated that it is often misleading[3,5]), it nonetheless plays an important role in astrophysics. For example, pulsar electrodynamics proposes that magnetosphere particles are threaded by magnetic field lines and rotate rigidly with a star.[11] Here we present experimental and analytical evidence of why the concept of frozen-in field lines is often misleading.

It has been established that under certain conditions, plasma can freely cross magnetic field lines[6]. Here we develop and extend the viewpoint of Bellan.

II. ANALYSIS ON THE EFFECT OF FROZEN-IN FIELD LINES

A. contradiction in the theory of frozen-in field lines

Although the frozen-in field concept is the "bed-rock" concept underlying ideal MHD, it currently lacks a specific definition. The two definitions given below will be familiar to many readers (Discussion of multifarious definitions will be deferred to Table IV):

1. The magnetic flux through a conducting fluid is conserved (we designate this definition Theorem 1).
2. Any motion of conducting fluid, perpendicular to the field lines, carries the field lines with the fluid. In other words, a highly conducting fluid cannot freely traverse magnetic field lines (we designate this definition Theorem 2)

The frozen-in field concept follows from the frozen-in field equation

$$\frac{\partial \boldsymbol{B}}{\partial t} = \nabla \times (\boldsymbol{v} \times \boldsymbol{B}), \tag{1}$$

where $\boldsymbol{v}$ is the relative velocity of the conductive fluid element moving through the magnetic field, and $\boldsymbol{v}\times\boldsymbol{B}$ is the motional electromotive field.

Theorem 1 follows unequivocally from the frozen-in equation (1). Theorem 2, on the contrary, conflicts with the frozen-in equation. We prove this statement as follows: If Theorem 2 holds (i.e. the motion of conducting fluid, perpendicular to the field lines, vanishes anywhere at any time), then $\boldsymbol{v}\times\boldsymbol{B}=0$ must be true. However, if $\boldsymbol{v}\times\boldsymbol{B}=0$ is always true, the frozen-in equation (1) becomes trivialized. Hence, Theorem 2 conflicts with the frozen-in equation. This may be restated in another way. Equation (1) describes a (time-varying) disturbance of magnetic field which depends on the traverse motion of fluid relative to the field lines. That is, the frozen-in equation (1) essentially describes the relationship between the magnetic disturbance and the traverse motion of fluid (Therefore, we shall rename Equation (1) as the "traverse-disturbance equation" to emphasize the indispensability of traverse motion). If Theorem 2 always holds and the traverse motion is absolutely debarred, the traverse-disturbance equation becomes apparently redundant (reducing to the trivial case 0 = 0). In other words, from an equation which describes the relationship between A and B, such a result that the B does not exist originally should not be obtained. Unluckily, Theorem 2 is actually come from such erroneous logic.

From Equation (1), it follows that the magnetic flux within a conducting fluid is always conserved. However,

---

[a)] Electronic mail: zx.liang55@gmail.com

magnetic flux can also be conserved in the following scenarios:

1. All field lines within a conducting fluid move along with the conducting fluid element.
2. The magnetic field lines entering a given fluid element are equal to those leaving the element at all times.

The second item describes the general condition of magnetic flux conservation, while the first describes a special case of item 2. If the magnetic field lines entering a given fluid element are equal to those leaving the element and are zero at all times, item 2 reduces to item 1. Theorem 2 of the frozen field concept applies strictly to the special condition, and ignores the general condition, thereby rendering the concept frequently misleading.

B. conditions for freely traversing magnetic field

Following discussion regarding the conflict between Theorem 2 and the traverse-disturbance equation, we discuss the conditions under which the conducting fluid can freely traverse magnetic field without magnetic disturbance and magnetic resistance.

From Equation (1), we realize that

1. In an ideal conducting fluid, the magnetic disturbance can arise only from the cross motion of conducting fluid. In the case of no cross motion, there can be no magnetic disturbance.
2. In the event that $\boldsymbol{v}\times\boldsymbol{B}\neq 0$, provided that $\nabla\times(\boldsymbol{v}\times\boldsymbol{B})=0$, magnetic disturbance must be absent.

This indicates that fluid cross motion is a necessary, but not a sufficient, condition for magnetic disturbance. The magnetic disturbance arising from the motion of conducting fluid is co-existent with magnetic resistance; neither phenomenon can appear by itself, since both arise from one and the same eddy current. Therefore, the cross motion is also a necessary but not sufficient condition for magnetic resistance. Since magnetic disturbance and magnetic resistance are the sole characteristics of magnetic-frozen states, we posit that the cross motion is a necessary but not sufficient condition for the magnetic-frozen states. In other words, even if a conducting fluid crosses magnetic field lines, the magnetic-frozen states can be absent provided that the special conditions $\boldsymbol{v}\times\boldsymbol{B}\neq 0$ and $\nabla\times(\boldsymbol{v}\times\boldsymbol{B})=0$ are satisfied.

It is known that the essential condition for a force acting between the fluid and the magnetic field is the presence of an eddy current within the conductive fluid, while the essential condition for an eddy current is presence of an eddy motional electromotive field $\boldsymbol{v}\times\boldsymbol{B}$. When both sides of Equation (1) are zero, the motional electromotive field within the fluid is irrotational. In this case, neither an eddy electromotive field nor an eddy current in the conductive fluid exist. Consequently, no magnetic force is exerted on the fluid when it crosses a magnetic field with velocity $\boldsymbol{v}$.

To determine the conditions under which the fluid can freely traverse magnetic field without magnetic disturbance and magnetic resistance, we expand Equation (1) as

$$\frac{\partial \boldsymbol{B}}{\partial t}=\boldsymbol{v}(\nabla\cdot\boldsymbol{B})-\boldsymbol{B}(\nabla\cdot\boldsymbol{v})-(\boldsymbol{v}\cdot\nabla)\boldsymbol{B}+(\boldsymbol{B}\cdot\nabla)\boldsymbol{v}. \quad (2)$$

The first term on the right-hand side involves the divergence of the magnetic field $\nabla\cdot\boldsymbol{B}$, which, according to Maxwell's equation, is zero always.

The factor $\nabla\cdot\boldsymbol{v}$ in the second term is the divergence of the velocity field. If the density distribution of the fluid remains constant, this term also equates to zero.

When each conductive fluid element moves along a path of constant magnetic field $\boldsymbol{B}$ (namely, an isomagnetic surface), the third term $(\boldsymbol{v}\cdot\nabla)\boldsymbol{B}$ additionally vanishes.

If the fluid elements that synchronously cross the same magnetic field line have the same velocity (i.e., do not exhibit differential movement), the fourth term $(\boldsymbol{B}\cdot\nabla)\boldsymbol{v}$ vanishes also. In general, the differential movement is related to the shape change of the fluid.

Equation (2) incorporates three essential factors that control the interaction between a conductive fluid and a magnetic field:

1. The expansion or compression of the fluid, i.e., the change in density;
2. The differential movement of the fluid, i.e., the shape change of the fluid; and
3. The change in magnetic field along the path of movement.

These essential factors indicate that the frozen-in phenomenon is related to distinct changes. Without these changes, the magnetic-frozen phenomenon cannot develop.

If a fluid of unchanging density distribution and shape can be regarded as a quasi-rigid body, the following simple law applies: *When crossing a magnetic field along an isomagnetic surface, a rigid or quasi-rigid body will not be subject to a magnetic resistance and the frozen-in phenomenon will not occur.* This law fits the scenarios shown in Figures 1, 2 and 3. The three scenarios share two characteristics:

1. The motional electromotive field $\boldsymbol{v}\times\boldsymbol{B}$ inside the fluid is nonzero and curl-free everywhere.
2. All of the fluid elements cross the magnetic field along an isomagnetic surface.

In the cases of Figs. 1 and 2, although the motional electromotive field $\boldsymbol{v}\times\boldsymbol{B}$ inside the fluid is irrotational, a rotational field $\boldsymbol{v}\times\boldsymbol{B}$ exists at the boundary of the conductive body. However, $\nabla\times(\boldsymbol{v}\times\boldsymbol{B})\neq 0$ is only a necessary, but not a sufficient, condition for the appearance of eddy currents. The eddy plane at the boundary is always perpendicular to the boundary surfaces; furthermore, no conducting matter exists outside the boundary and eddy currents cannot cross the boundary surfaces. This ensures that no eddy current can appear at the boundary and hence the magnetic field cannot be modified. In the case of Figure 3, the conductive body is a ring, and there has neither front boundary nor rear boundary in the direction of movement. The boundary analysis is appropriate only for the top and bottom surfaces of the conductive body.

For the case shown in Figure 3, even if the outer and inner fluids rotate at different angular speeds, i.e. $\omega = f(r)$, no magnetic resistance acts on the fluid. However, if the differential rotation can be described as $\omega = f(z)$ (where z is the axial coordinate), a magnetic resistance will act on the fluid. This distinction arises from the fourth term of the right-hand side of Equation (2), $(\boldsymbol{B}\cdot\nabla)\boldsymbol{v}$. If $\omega = f(r)$, then $(\boldsymbol{B}\cdot\nabla)\boldsymbol{v} = 0$ and no magnetic force is felt; if $\omega = f(z)$, then $(\boldsymbol{B}\cdot\nabla)\boldsymbol{v}\neq 0$ and a magnetic resistance is felt. This result is consistent with the law of iso-rotation (Ferraro's theorem), which dictates that in the steady state, angular velocity is constant along magnetic lines[4]. However, if $\omega = f(r)$, the angular velocity varies only along the direction perpendicular to the field lines.

Alfvén & Fälthammar[4] has described an example similar to that in Figure 1, for which the conductive fluid can cross a uniform magnetic field under certain conditions. Our work extends this idea by softening the constraint conditions for other cases of non-uniform magnetic field and/or differential rotation of the conductive fluids.

Because the case shown in Figure 3 illustrates the situation in the vicinity of the equatorial plane of an aligned pulsar, we come to an important conclusion: the plasma near the equatorial plane of an aligned pulsar can cross the magnetic field and cannot be frozen with the magnetic field lines; in other words, the plasma can neither drive the magnetic field lines, nor be driven by the magnetic field lines. This conclusion is supported by the results of our magnetohydrodynamic experiments (see next section).

Figures 1, 2, and 3 do not include all of the scenarios in which fluid can freely cross magnetic field lines. The conditions in these three scenarios merely ensure that all terms of the right-hand side of Equation (2) are zero. More generally, even if some terms are non-zero, so long as the terms sum to zero, the fluid can freely cross the magnetic field lines. For example, if a conducting fluid element undergoes an apropos expansion rate, it can cross magnetic field lines against the direction of magnetic gradient.

Figure 4 shows a scenario in which the fluids experience magnetic resistance resulting in magnetic disturbance. The significance of isomagnetic surfaces is highlighted by comparing Figures 2 and 4.



FIG. 1. The first non-frozen scenario. Moving across a uniform magnetic field, an ideal conductive fluid neither experiences a frozen resistance nor carries the magnetic field lines, contrary to the expectations of some plasma scientists. Here, the distributions of the charges and electric fields have been simplified.



FIG. 2. The second non-frozen scenario. For a conductive fluid moving across a non-uniform magnetic field, provided that the fluid elements move along an isomagnetic surface, there is no frozen-in effect.

If the magnetic field $\boldsymbol{B}$ changes in the direction of fluid movement, the fluid will generally experience resistance. In particular, if the length of fluid along the flow direction is greater than the extent of the magnetic field, as shown in Figure 5, magnetic resistance is inevitable at the boundary of the magnetic field because the magnetic field changes sharply there.

It is known that a conductive quasi-rigid body crossing



FIG. 3. The third non-frozen scenario. In a cylindrically symmetric magnetic field, rotation of a conductive fluid around a magnetic axis does not result in the frozen-in phenomenon, even if the outer and inner fluids rotate at different angular speeds. This scenario mirrors the situation in the vicinity of the equatorial plane of an aligned pulsar.

a uniform magnetic field is not subject to magnetic force. Generally, a steady magnetic field can be regarded as a superposition of uniform and non-uniform components. Only the non-uniform component can exert a resistance on the fluid, as shown in Figure 4. However, the non-uniform component does not always mount a resistance, as we have shown in Figures 2 and 3. Consequently, in plasma physics, the frozen-in effect should be related to the magnetic gradient in the direction of fluid motion, but not to the uniform magnetic field component. From this perspective, the frozen-in concept is consistent with standard electromagnetism, and is no longer misleading.

C. effects of the polarization charges and electrostatic field



FIG. 4. A magnetically frozen scenario. If the fluid velocity has a component in the direction of the magnetic field gradient, movement of the fluid will result in the frozen-in effect.



FIG. 5. Another magnetically frozen scenario. If the extent of the magnetic field is smaller than that of the fluid, even if the magnetic field is uniform, a frozen-in effect will occur because of the high magnetic field gradient near the magnetic field boundary.

According to conventional physics, a straight wire, closed wire or conducting plate (see Figure 1) moving perpendicularly in a uniform magnetic field can neither produce electric current nor experience resistance from the magnetic field. Thus, the conductors cannot be frozen within the magnetic field lines. Because this phenomenon is independent of type or conductivity of the materials, even if the objects shown in Figure 1 comprise a conductive fluid such as mercury or plasma, the frozen-in effect does not apply. However, in the context of plasma physics, if the fluids in Figure 1 are conductive, it may be considered that they will be frozen in the magnetic field and cannot pass freely through field lines. This presents an apparent contradiction between conventional physics and plasma physics.

The ambiguity of the concept of frozen-in field lines arises from an oversight. In previous studies of the frozen-in effect, the effects of the polarization charges and electrostatic field $\boldsymbol{E}$ shown in Figures 1, 2 and 3 have been disregarded. In fact, it is the electric field $\boldsymbol{E}$ that produces a fluid drift and "thaws" the magnetic-frozen states. A simple calculation using the drift theory of charged particles[7] will verify that the drift velocity of both the ions and electrons produced by $\boldsymbol{E}$ and $\boldsymbol{B}$ is always equal to the macroscopic moving velocity of the fluid, i.e., $\boldsymbol{v}_{\text{drift}} \equiv \boldsymbol{v}$. The direction of the drift velocity $\boldsymbol{v}_{\text{drift}}$ is always perpendicular to both the electric and magnetic fields. Of course, to ensure that $\boldsymbol{v}_{\text{drift}} \equiv \boldsymbol{v}$, no eddy currents are permitted inside the fluid (as is true for Figures 1, 2 and 3).

Bellan has demonstrated[6] that a plasma immersed in a static magnetic field $\boldsymbol{B}$ and a static electric field $\boldsymbol{E}$, with $\boldsymbol{E}$ perpendicular to $\boldsymbol{B}$, will freely drift across the magnetic field lines. On the basis of Bellan's viewpoint, we claim that the static electric field $\boldsymbol{E}$, which is established by polarization charges, can naturally appear under certain conditions. The process by which this can occur is summarized below:

A conducting fluid moving across the magnetic field lines induces an irrotational motional electromotive field $\boldsymbol{v}\times\boldsymbol{B}$ inside the fluid; → As a result of this irrotational motional electromotive field, polarization charges appear inside the fluid; → The polarization charges result in a static polarization electric field $\boldsymbol{E} = -\boldsymbol{v} \times \boldsymbol{B}$; → which combines with the magnetic field $\boldsymbol{B}$ to "drive" the fluid at drift velocity $\boldsymbol{v}_{\rm drift}$ equal to $\boldsymbol{v}$.

Finally, the inertial velocity of the moving fluid is the same as that in magnetic-field-free space.

Both the eddy current and the polarization charges are essentially the electric phenomena arising from the crossing motion of conducting fluid, and jointly influence the interaction between the fluid and the magnetic field. However, in previous reports of the concept of frozen-in field lines, the effect of the eddy current is always considered, while the effect of the polarization charges remains overlooked. This issue deserves serious attention.

When an individual charged particle travels through a magnetic field, it is subject to Lorentz forces which compel it to gyrate around the magnetic field lines. However, within a conductive fluid, the presence of polarization charges and an electric field create a very different situation. Under some special conditions, such as those illustrated in Figures 1, 2, and 3, the static electric field produced by the polarization charges can counteract the Lorentz force and the fluid elements will cross the magnetic field without resistance.

To study the interaction between a conductive fluid and a magnetic field, both the curl and divergence of the motional electromotive field should be considered. The curl affects the eddy current while the divergence affects the polarization charges and electric field. Although we can create situations in which the magnetic field is curl-free, such as those illustrated in Figures 1, 2 and 3, the motional electromotive field $\boldsymbol{v} \times \boldsymbol{B}$ cannot be made divergence-free inside a fluid. Therefore, the divergence affects the fluid movement more generally than does the curl. However, because Equation (1), from which the frozen-in concept derives, contains no divergence term, the frozen-in concept cannot entirely and accurately describe the interaction between a conductive fluid and a magnetic field. This limitation is responsible for the situations in which the frozen-in concept is often misleading, as emphasized by Alfvén.[5]

D. movement of magnetic field lines

In the case of Figure 1, although magnetic disturbances are absent, it may be considered that the magnetic field lines are nonetheless carried by the fluid. However, this viewpoint would result in contradiction, as shown in Figure 6. The magnetic field in Figure 6 is initially uniform, and the quasi-rigid fluids B and A are respectively immobile and traveling. Because the fluid B is immobile, the magnetic field lines through that fluid must be immobile also. If the magnetic field lines through the fluid A are carried by fluid A (as shown in Figure 6c), regardless of reference frame, a magnetic disturbance must appear in the space between the fluids. However, foregoing analysis has indicated that a quasi-rigid fluid traveling within a uniform magnetic field cannot induce magnetic disturbance. Therefore, the magnetic field of Figure 6c is unrealistic and the viewpoint that the fluid always carries magnetic field lines must be discarded. The immobile fluid B ensures that magnetic disturbance appears in any reference frame, thereby precluding the use of the moving frame (which complicates the problem). The same effect is achieved if the velocities of the fluids in Figure 1 are assumed different.



FIG. 6. Different model leading to different corollaries. (a) The conductive fluid A moves across the uniform magnetic field. (b) Assuming that the fluid can traverse the magnetic field, the magnetic field will retain uniformity. (c) Assuming that the fluid can carry the magnetic field lines, the magnetic field will become non-uniform.

Our sentiments are embodied in the following quote

TABLE I. Contrast between two cases of the traverse-disturbance equation

	case A	case B
both sides of Equation (1)	$= 0$	$\neq 0$
property of magnetic field	steady magnetic field	time-varying magnetic field
eddy current in fluid	absence	presence
magnetic resistance	absence	presence
frozen-in phenomena	absence	presence
illustration	Figures 1, 2 and 3	Figures 4 and 5

from Feynman et al.[10]: "It makes no sense to say something like: When I move a magnet, it takes its field with it, so the lines of $\boldsymbol{B}$ are also moved. There is no way to make sense, in general, out of the idea of 'the speed of a moving field line.'" A magnetic field has essentially no movement attributes. Time-varying magnetic fields exist, whereas moving ones do not. While it can be intuitive and useful to regard magnetic field lines as moving, this viewpoint can lead to erroneous interpretation. Such a misleading concept may underlie Theorem 2 in the concept of frozen-in field lines.

E. two cases of the traverse-disturbance equation

Table I contrasts two cases of the traverse-disturbance equation and clarifies the relationship between the frozen-in phenomena and the traverse-disturbance equation.

III. EXPERIMENTS ON THE EFFECT OF FROZEN-IN FIELD LINES

To verify the above analysis, we conducted a series of magnetohydrodynamic experiments. The configuration is the same as that of Figure 3. For simplicity and clarity, the experiments were recorded and the videos of these can be viewed on YouTube. Figures 7 and 8 depict the device configurations and the sizes of the main components. Apart from the magnet, all components are made of non-magnetic material (plastic or copper). The mercury was placed in a cylindrical trough surrounding a rotating platform. At the magnet surface, the magnetic field intensity is about 0.6 T. The rotational rates are shown in the relevant videos.

The experiments were divided into two groups. The first group comprised three pairs of experiments to determine the conditions under which the magnet can drive rotation of the mercury. In each experiment the basis of comparison is an *aligned magnetic rotator*, a cylindrically symmetric magnet whose magnetic axis aligns with its rotation axis. The three experiments are as follows:

1. *Drive experiment A*[17] revealed the contrast between the aligned rotator and a rotator with magnetic axis orthogonal to the rotation axis. This contrast test indicates that the transverse magnetic field can drive the mercury to rotate, but the aligned magnetic field cannot do that.
2. *Drive experiment B*[18] revealed the contrast between the aligned rotator and a magnet with magnetic axis parallel to, but displaced from the rotation axis. This contrast test indicates that the eccentricity magnetic field can drive the mercury, but the coaxial magnetic field cannot do that.
3. *Drive experiment C*[19] revealed the contrast between the aligned rotator and a magnet that is not cylindrically symmetric. This contrast test indicates that the cylindrically asymmetric magnetic field can drive the mercury, but the cylindrically symmetric magnetic field cannot do that.

The first set of experiments demonstrated that an aligned magnetic rotator cannot drive rotation of the mercury. However, this does not necessarily imply the absence of a frozen magnetic effect in such cases, because the same effect will arise if the magnetic field is not co-rotating with the aligned magnet. The value of these experiments lies in their potential to resolve, at least partially, the following issue: Can an aligned pulsar drive its plasma and achieve the co-rotation state?



FIG. 7. Schematic figure of drive experimental device. The unit of length is millimeter (mm). The belt drives the rotation of the platform and magnet; the other components are stationary.

TABLE II. Summary of the analysis results

	magnetic frozen (column A)	magnetic "thaw"(column B)
induction phenomena	inductive current	polarized charges
induction electric field parameter	curl of electric field	divergence of electric field
correlative magnetic field parameter	magnetic field gradient	magnetic field strength
direction of fluid movement	parallel to magnetic gradient	perpendicular to magnetic gradient
component of the magnetic field	non-uniform	uniform
effect of magnetic force	resisting the crossing motion	impelling the crossing motion
illustration	Figures 4 and 5	Figures 1, 2 and 3



FIG. 8. Schematic figure of braking experimental device. The axis, support block and the magnet are stationary. The belt drives the rotation of the other parts. The unaccounted-for dimensions are the same as those of Fig. 7.

In the second group of experiments, the conditions under which rotating mercury can be slowed by a stationary magnet were determined. The mercury trough (rather than the central magnet) was spun until the mercury revolution reached a steady state. In each experiment the basis of comparison was an *aligned magnet*, a cylindrically symmetric magnet whose magnetic axis matched that of the mercury rotation axis. In this case, the mercury eventually revolved at the same rate as the trough. For the three comparison cases, braking forces reduced the rotation of the mercury. We used the same magnets in every pair of braking experiments, designated A, B and C and described below.

1. *Braking experiment A*[20] revealed the contrast between the aligned magnet and the orthogonal magnet. This contrast test indicates that the orthogonal magnetic field can slow down the mercury, but the aligned magnetic field cannot do that.

2. *Braking experiment B*[21] revealed the contrast between the aligned magnet and the off-axis magnet. This contrast test indicates that the eccentricity magnetic field can slow down the mercury, but the aligned magnetic field cannot do that.

3. *Braking experiment C*[22] revealed the contrast between the aligned magnet and the cylindrically asymmetric magnet. This contrast test indicates that the cylindrically asymmetric magnetic field can slow down the mercury, but the cylindrically symmetric magnetic field cannot do that.

The braking experiments demonstrated that a coaxial, cylindrically symmetric magnetic field[23] exerts no braking effect on the mercury. In other words, under the conditions of Figure 3, even with no radial electric field applied beforehand, the mercury can traverse the magnetic field lines without experiencing a braking force. The magnet can exert a braking effect on the mercury only when its magnetic axis is unaligned with the mercury rotation axis or when the magnetic field is not cylindrically symmetrical. This set of experiments also demonstrated that, even if the magnetic field in non-uniform, conductive fluid moving in the field experiences no frozen-in effect, so long as all of the fluid elements move along the isomagnetic surface. Thereby, the analyses in section II have been corroborated by our experimental results.

IV. DISCUSSION

For clarity, the analysis results are summarized in Table II. The concept of frozen-in field lines is in fact related to each entry in this table. Whether a magnetic field can "drive" or resist the crossing motion of fluid strongly depends on the distribution of the magnetic field and the direction of fluid movement. In general cases, both fluid and magnetic field should exist in a semi-frozen state. However, former studies have mainly focused on those properties listed under column A of Table II and have overlooked all of the column B aspects; that is, the frozen-in effect has been emphasized while the drift effect has been largely ignored. Consequently, the concept of frozen-in field lines has often been misleadingly stated as *a conductive fluid cannot freely cross a magnetic field.*

Table III presents a summary of the experimental results. Here, it is emphasized that no exchange of angular momentum occurs between the magnet and the mercury under conditions in which the fluid elements move along the isomagnetic surface. The extents of axial alignment and symmetry (essentially, the magnetic gradient along the path of movement), thus control the exchange of angular momentum. Naturally, a change in the magnetic

TABLE III. Summary of the experimental results

	Driving experiment results	Braking experiment results
aligned axis and cylindrical symmetry cases	no[a]	no
orthogonal magnetic field	yes[b]	yes
off-axis magnetic field	yes	yes
cylindrically asymmetric magnetic field	yes	yes

[a] "no" implies the absence of magnetic frozen phenomenon.
[b] "yes" implies the presence of magnetic frozen phenomenon.

TABLE IV. Summary of the expressions of frozen-in field concept

correct expressions	incorrect expressions
Any two elements of the fluid that are at one instant on a common magnetic field line will be on a common magnetic field line at any other instant.[9]	Any element of the fluid that is at one instant on a magnetic field line will be on the original magnetic field line at any other instant.
A highly conducting fluid generally cannot freely traverse magnetic field lines along the direction of magnetic gradient.	A highly conducting fluid cannot freely traverse magnetic field lines.
The magnetic flux in any conducting fluid element is conserved.	The magnetic lines in any conducting fluid element will move along with the fluid element.

flux passing through the fluid elements is a necessary condition for angular momentum exchange.

From the experimental and analytical results presented here, we conclude that when a conducting fluid moves in a magnetic field, whether or not resistance acts on the conducting fluid depends solely on the spatial variation of the magnetic field ($\Delta \boldsymbol{B}$), and is not related to the magnetic field intensity ($\boldsymbol{B}$). This corollary may alleviate some of the current misunderstanding of the concept of frozen-in field lines. For many astrophysical objects, the magnetic field may be very strong, but the magnetic field gradient in the direction of plasma movement is very small, especially for cases similar to that shown in Figure 3. Consequently, the magnetic frozen effect is much weaker than expected and the plasma can freely cross the magnetic field lines. In other words, in many astrophysical scenarios, the magnetic frozen effect can be ignored provided that the magnetic gradient along the path of plasma movement can be ignored.

In fact, Alfvén's warning that the concept of frozen-in field lines is often misleading has not been taken seriously by astrophysical scientists, especially by pulsar scientists. To date, almost all of the papers addressing the topic of pulsar radiation have been misled by the frozen-in field concept. These papers unanimously assume that the particles of magnetosphere can move freely along magnetic field lines but cannot freely traverse them.

The concept of frozen-in field lines can be expressed in several forms, which are summarized in Table IV. All of the expressions listed in the right column of table IV are derived from Theorem 2, and are hence incorrect or incomplete.

V. SUMMARY

1. We suggest renaming the frozen-in equation as the traverse-disturbance equation, to emphasize the universality of traverse motion and to reduce opportunities for erroneous interpretations arising from the concept of frozen-in field lines.
2. We suggest emphasizing the especial significance of isomagnetic surfaces and polarization charges, in order to entirely and accurately understand the interaction between a conductive fluid and a magnetic field.
3. Whether a quasi-rigid conducting fluid can freely traverse magnetic field lines or not depends solely on the gradient of the magnetic field in the direction of fluid movement and is independent of the magnetic field strength.
4. Because Theorem 2 conflicts with the traverse-disturbance equation, we strongly recommend that it be discarded.

[1] S.-I. Akasofu, *Space Sci. Rev.* **21**, 489 (1978).
[2] H. Alfvén, *Nature* **150**, 405 (1942).
[3] H. Alfvén, *Tellus* **10**, 104 (1958).
[4] H. Alfvén and C.-C. Fälthammar, *Cosmical electrodynamics, fundamental principles,* 2nd ed. (Clarendon Press, 1963), p. 110, p.190 Fig. 5.10.
[5] H. Alfvén, *J. Geophys. Res.* **81**, 4019 (1976).
[6] P. M. Bellan, *Fundamentals of plasma physics,* (Cambridge University Press 2006), p. 100.
[7] J. A. Bittencourt, *Fundamentals of plasma physics,* 3rd ed. (Springer 2004), p. 312, p. 51.
[8] T. J. M. Boyd and J. J. Sanderson, *The physics of plasmas.* (Cambridge University Press 2003), p. 79.
[9] C.-G. Fälthammar,. *Am. J. Phys.* **74**, 454 (2006).

[10] R. P. Feynman, R. B. Leighton and M.Sands, *The Feynman Lectures on Physics*, Vol. 2 (Addison wesley 1964), p. 13-10.

[11] P. Goldreich and W. H. Julian, *Astrophys. J.* **157**, 869 (1969).

[12] I. A. Ivonin, V. P. Pavlenko, and H. Persson, *Phys. Plasmas* **5**, 2893 (1998).

[13] R. Lundin, M. Yamauchi, J.-A. Sauvaud, and A. Balogh, *Annales Geophysicae* **23**, 2565 (2005).

[14] H. Nishida and T. Abe, *Phys. Plasmas* **17**, 052901 (2010).

[15] T. Passot, V. Ruban, and P. L. Sulem, *Phys. Plasmas* **13**, 102310 (2006).

[16] R. L. Stenzel, J. M. Urrutia, and M. C. Griskey, *Phys. Plasmas* **6**, 4458 (1999).

[17] Z. X. Liang, and Y. Liang, http://www.youtube.com/watch?v=YCc4ybfKiY0, YouTube (2009).

[18] Z. X. Liang, and Y. Liang, http://www.youtube.com/watch?v=ZiVNxVAqUKc, YouTube (2009).

[19] Z. X. Liang, and Y. Liang, http://www.youtube.com/watch?v=SBAzUBzc2xM, YouTube (2009).

[20] Z. X. Liang, and Y. Liang, http://www.youtube.com/watch?v=C7DxRoVdtxw, YouTube (2009).

[21] Z. X. Liang, and Y. Liang, http://www.youtube.com/watch?v=p9UzfT6UHcE, YouTube (2009).

[22] Z. X. Liang, and Y. Liang, http://www.youtube.com/watch?v=Eq0B5XR_o8U, YouTube (2009).

[23] Many discussions in this paper assume cylindrical symmetry. In the experiments, however, although the magnetic field generated by the cylindrical magnet does not completely meet the condition of cylindrical symmetry, the center of the magnetic field is almost perfectly cylindrically symmetric. Therefore, most of our discussions are limited to the middle of the magnetic field.